




American Physical Therapy Association

OPTIMAL Difficulty-Follow-Up

Instructions: Please circle the level of difficulty you have for each activity today.	Able to do without any difficulty	Able to do with little difficulty	Able to do with moderate difficulty	Able to do with much difficulty	Unable to do	Not applicable
1. Lying flat	1	2	3	4	5	9
2. Rolling over	1	2	3	4	5	9
3. Moving-lying to sitting	1	2	3	4	5	9
4. Sitting	1	2	3	4	5	9
5. Squatting	1	2	3	4	5	9
6. Bending/stooping	1	2	3	4	5	9
7. Balancing	1	2	3	4	5	9
8. Kneeling	1	2	3	4	5	9
9. Walking-short distance	1	2	3	4	5	9
10. Walking-long distance	1	2	3	4	5	9
11. Walking-outdoors	1	2	3	4	5	9
12. Climbing stairs	1	2	3	4	5	9
13. Hopping	1	2	3	4	5	9
14. Jumping	1	2	3	4	5	9
15. Running	1	2	3	4	5	9
16. Pushing	1	2	3	4	5	9
17. Pulling	1	2	3	4	5	9
18. Reaching	1	2	3	4	5	9
19. Grasping	1	2	3	4	5	9
20. Lifting	1	2	3	4	5	9
21. Carrying	1	2	3	4	5	9

22. Thinking about all of the activities you would like to do, please mark an "X" at the point on the line that best describes your overall level of difficulty with these activities today.



© 2005, 2006 American Physical Therapy Association. All rights reserved. No part of this instrument may be reproduced, stored in a retrieval system, or transmitted in any form or by any means, electronic, mechanical, photocopying, or otherwise without prior permission of the American Physical Therapy Association. Contact permissions@apta.org or visit www.apta.org/publications.

Adapted/revised in July 2005 and August 2006 with permission of APTA from Guccione AA, Mielenz TJ, De Vellis RF, et al. Development and testing of a self-report instrument to measure actions: Outpatient Physical Therapy Improvement in Movement Assessment Log (OPTIMAL). *Phys Ther.* 2005;85:515-530.